

Interaction of secreted aspartic proteinase *Candida albicans* with ZnCl₂: Complex formation and catalytic activity

Kutyreva M., Ulakhovich N., Gataulina A., Mukhametzyanova A., Khaldeeva E.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The aim of work is to estimate ion Zn(II) influence on catalytic activity of secreted aspartic proteinase *Candida albicans* (SAP2). System ZnCl₂ - SAP2 showed complex formation. Composition of [ZnmSAP2n] complex was 1:1, the stability constant was 4.73 ± 0.20 . Affinity constants in system SAP2 - human serum albumin (HSA) - ZnCl₂ were determined at Skatchard coordinates. The SAP2 was found to have a one section for linking with the substrate and one section for linking with modulator. The affinity constants in SAP2 - HSA system decreased in presence of ZnCl₂. Complex forming of SAP2 - ZnCl₂ system resulted in reduction of enzyme-substrate bonding. Proteolytic activity of SAP2 towards HSA at presence ZnCl₂ as modulator was estimated. The inhibition effect in range 1×10^{-4} - 4×10^{-6} and 6×10^{-7} - 1×10^{-8} M was observed. The SAP2 activation effect catalyzed by ZnCl₂ at 5×10^{-7} - 5×10^{-6} M concentrations were fixed for the first time. © IDOSI Publications, 2013.

<http://dx.doi.org/10.5829/idosi.wasj.2013.27.09.13747>

Keywords

Enzyme activity, Metal complex, Proteinase *Candida albicans*, Zinc ion